

# Effects of Eccentric Exercise on Muscle Architecture in Adults: A Systematic Review

## Efectos del Ejercicio Excéntrico en la Arquitectura Muscular en Adultos: Una Revisión Sistemática

Lizama-Pérez, R.<sup>1,2</sup>; Chiroso-Rios, I.<sup>1</sup>; Chiroso-Rios, L.<sup>1</sup>; Olave, E.<sup>3</sup>; Ferragut, C.<sup>4</sup>; Vila, H.<sup>5</sup> & Jerez-Mayorga, D.<sup>6</sup>

**LIZAMA-PÉREZ, R.; CHIROSA-RIOS, I.; CHIROSA-RIOS, L.; OLAVE, E.; FERRAGUT, C.; VILA, H. & JEREZ-MAYORGA, D.** Effects of eccentric exercise on muscle architecture in adults: A systematic review. *Int. J. Morphol.*, 40(2):425-432, 2022.

**SUMMARY:** The purpose of this systematic review was to determine the effects of eccentric training on muscle architecture in the adult population. Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) statements were followed using keywords associated with architecture muscular and eccentric training. Four databases were used: PubMed, Scopus, SPORTDiscus and Web of Science. Methodological quality was assessed using the PEDro scale. A total of 1260 articles were retrieved, 18 included in this review. The parameters most frequently evaluated in the studies consulted were pennation angle (PA), fascicle length (FL), and muscle thickness (MT). These were assessed mainly in lower limb muscles such as biceps femoris long head (BF<sub>lh</sub>), vastus lateralis (VL), medial gastrocnemius (MG) and lateral gastrocnemius (LG), respectively. Eccentric training for at least four weeks generates adaptations in these parameters, mainly by increasing MT with FL and decreasing PA, determining muscle function. These results provide evidence on the effects of eccentric training on muscle architecture, which could be helpful to prevent injuries and favor muscle recovery processes.

**KEY WORDS:** Muscular architecture; Eccentric training; Muscle power; Pennation angle; Muscle thickness; Fascicle length; Muscle quality.

## INTRODUCTION

To understand in greater depth the situation of skeletal muscle in sports or injury rehabilitation contexts, it is necessary to know the characteristics of muscle tissue in a broader and more functional sense (Suchomel *et al.*, 2016). One of the most objective and global concepts that contemplate this tissue's physiological and functional capacity is muscle quality (MQ) (Fragala *et al.*, 2014). Muscle quality is composed of four dimensions (muscle composition, architecture, ultrastructure, and functional unit) and two indexes (relative strength and muscle quality index) with architecture being one of the least explored factors (Fragala *et al.*, 2015). Its evaluation can provide us with the muscle's capacity to generate strength, power, or functionality (Jerez-Mayorga *et al.*, 2020).

Muscle architecture (MA) is defined as the arrangement of muscle fibers within a muscle about the axis of force generation, becoming one of the most determinant components of muscle function, which can be associated with functional and health components in individuals (Lieber & Fridén, 2000; Lieber & Ward, 2011; Naimo *et al.*, 2021). Therefore, MA is a fundamental element to be considered in assessing MQ (Naimo *et al.*). There is variability in the architecture of a muscle; however, generally, two types of architectural arrangements are described, longitudinal muscles (muscle fibers are arranged parallel to the force-generating axis) and pennate muscles (fibers are oriented at one or more angles concerning the force-generating axis) (Lieber & Fridén; Lieber & Ward). Several parameters can

<sup>1</sup> Department Physical Education and Sports, Faculty of Sport Sciences, University of Granada, 18011 Granada, Spain.

<sup>2</sup> Departamento de Ciencias Morfológicas, Facultad de Medicina y Ciencia, Universidad San Sebastián, General Lagos 1163, Valdivia 509000, Chile.

<sup>3</sup> Facultad de Medicina, Universidad de La Frontera, Temuco, Chile.

<sup>4</sup> Faculty of Medicine and Health Sciences, University of Alcalá, 28871 Alcalá de Henares, Spain.

<sup>5</sup> Faculty of Education, University of Vigo, 36905 Pontevedra, Spain.

<sup>6</sup> Exercise and Rehabilitation Sciences Laboratory, School of Physical Therapy, Faculty of Rehabilitation Sciences, Universidad Andres Bello, Santiago, Chile.

be considered; however, these are often conditioned by the assessment method used; these parameters include muscle thickness (MT); muscle length (ML), fascicle length (FL), pennation angle (PA), and physiological cross-sectional area (PSCA, the latter two being most closely related to muscle force generation (Lieber & Fridén; Lieber & Ward).

Understanding the architectural adaptations of skeletal muscle to different types of training has not yet been established by the scientific community. Several studies have attempted to determine the modifications of the architectural parameters in reference to different exercise programs; however, it is still not clear which is the best type of exercise to produce effective changes in these parameters (Narici *et al.*, 2016). Recently, Gerard *et al.* (2020), through a meta-analysis, determined the effects of eccentric exercise on the muscular architecture of the long head of the biceps femoris, concluding that it produces adaptations by increasing its MT, FL and decreasing the PA, as well as producing adaptations in the strength of the hamstring muscles. This could favor the recovery processes in the face of muscular injuries or even prevent them (Blazevich & Sharp, 2005).

However, the effects of eccentric training on other muscles have not been systematically reviewed. Therefore, this review aimed to assess the effects of eccentric training on muscle architecture in the adult population.

## MATERIAL AND METHOD

PRISMA guidelines were used (The Preferred Reporting Items for Systematic Reviews and Meta-Analyses). The protocol for this review was registered in INPLASY (2021120094).

**Search strategy.** The search was performed by two authors (RL-P and DJ-M). The databases used were Pubmed, Scopus, SPORTDiscus and Web of Science. The search was performed from inception until March 2021. The following keywords were included: “eccentric training”, “eccentric contraction”, “eccentric exercise”, “lengthening contraction”, “negative work”, “muscle architecture”, “pennation angle”, “fibre length”, “fiber length”, “fascicle length”, “cross-sectional area”, “muscle thickness”. The search was not limited in years.

**Inclusion criteria.** Articles that met the following criteria were included in this review: (I) subjects >18 years old, (II) Eccentric training program longer than four weeks (III) Studies with randomized clinical trial design, (IV) studies reporting measures of muscle architecture: “pennation

angle”, “fascicle length”, “muscle thickness”, (V) full text available, and (VI) articles in English. In addition, we excluded all those articles that (I) Eccentric training programs of less than four weeks (II) conference presentations, theses, books, editorials, review articles and expert opinions, (III) duplicate articles, and (IV) articles in which the principal or secondary authors did not respond to e-mail requests.

**Study selection.** The articles retrieved from the search were entered into the Rayyan QCRI application (Ouzzani *et al.*, 2016). This app assists the article selection process, optimizing review time and allowing collaborative work among researchers. (Available for free at <http://rayyan.qcri.org> (accessed on 27 March 2021)).

Duplicate articles were eliminated, and two investigators (RL-P and DJ-M) independently reviewed titles and abstracts to identify articles that met the eligibility criteria. A third investigator (LC-R) was consulted and resolved by consensus in case of discrepancies. Finally, the selected articles were read in total, and the reference list was reviewed for relevant articles that could be included.

**Data extraction.** An Excel template will be used for data extraction for each manuscript selected for review. The following information will be considered: author, year, aim, architectural parameter, sample size, age, population, physical activity level, number of participants, eccentric training protocol, results, and conclusions.

**Methodological quality.** The quality of the evidence of the articles included in this review was assessed using the PEDro scale, which is based on criteria that identify whether the RCTs have sufficient internal validity and statistical information to interpret the results (external validity (item 1), internal validity (items 2-9) and statistical reporting (items 10-11)). Each item is classified as yes or no (1 or 0) according to whether the study met the criterion. The total score is from item 2 to 11, so the maximum score is 10 (Cashin & McAuley, 2020). Two independent investigators (RL-P and DJ-M) evaluated the articles using this scale. In case of discrepancy, a third evaluator (LC-R) was consulted. About the quality of evidence, it has been suggested that scores < 4 are considered poor quality, 4 - 5 moderate, 6 - 8 good and 9 - 10 excellent (Cashin & McAuley).

## RESULTS

**Article Selection.** From the initial search, 1260 articles were retrieved, of which 726 were eliminated because they were duplicates. One additional article was identified from another

source. After evaluating titles and abstracts, 512 articles were excluded because they did not meet the inclusion criteria, leaving 23 articles for full-text analysis. All the assessed articles presented a control group in this systematic review.

Of the 23 articles, four were eliminated because they did not evaluate the results and one article did not evaluate the target population. Thus, 18 articles were selected, their reference lists were checked, and no new articles were found (Fig. 1). In addition, related studies were included in the drafting of the text.

**Study Characteristics.** In total, 506 subjects participated in the studies, with a mean of 29.7 of total participants for each study. The number of participants ranged from 12 to 49. The mean age of participants ranged from 18 to 29 years old with a mean of  $23.4 \pm 2.9$  years. Only one study did not mention the age of subjects (Mendiguchia *et al.*, 2020). Some participants engaged in recreational physical activity (Coratella *et al.*, 2015; Timmins *et al.*, 2016; Bourne *et al.*, 2017; Seymore *et al.*, 2017; Alonso-Fernandez *et al.*, 2020; Marusic *et al.*, 2020; Presland *et al.*, 2020) moderate physical activity (Ribeiro-Alvares *et al.*, 2018; Abián *et al.*, 2020), and others were physically active (Cadore *et al.*, 2014; Marzilger *et al.*; 2019, 2020; Mendiguchia *et al.*; Timmins *et al.*, 2021). In four studies, the subjects' physical activity

level is not mentioned (Guilhem *et al.*, 2013; Franchi *et al.*, 2014; Sanz-López *et al.*, 2016, 2017). The characteristics of the studies are summarized in Table I.

In this review, the evaluation of the methodological quality was assessed between the two reviewers in the totality of the articles (100 %). The results show that 89 % of the reviewed studies have a "Good" methodological quality, with PEDro 6-8 scale values. The results of this evaluation are shown in Table II.

**Effect of Eccentric Strength Training on muscle architecture.** Fifteen studies evaluated the effects of eccentric training on PA. Eleven studies found a decrease in PA; of these, in five (Mendiguchia *et al.*; Ribeiro-Alvares *et al.*; Marusic *et al.*; Timmins *et al.*, 2016, 2021), NHE strength training was performed in two (Sanz-Lopez *et al.*, 2016, 2017), eccentric overload training was conducted, in one (Presland *et al.*) eccentric training with flywheel was performed and in three others (Guilhem *et al.*; Franchi *et al.*; Marzilger *et al.*, 2020) isokinetic eccentric training. Four studies found an increase in PA, in one eccentric training with heel drop exercise (Alonso Fernández *et al.*), one eccentric resistance training with workload (Guilhem *et al.*), one declined squat training with one leg at two speeds (Abián *et al.*) and one isokinetic unilateral eccentric training (Coratella *et al.*)




Fig. 1. Flow Chart for the systematic review.

Table I. Characteristics of individual studies.

Study	Group	N	Age (Mean ± SD)	Activity Level
Alonso- Fernández <i>et al.</i> (2020)	Control	20	26.7 ± 2.8	Recreational physical Activity
	Eccentric training with heel drop exercise	25	25.2 ± 3.1	
Bourne <i>et al.</i> (2017)	Control	10	21.3±3.7	Recreational physical Activity
	Nordic hamstring exercise	10	21.6±3.2	
	Hip extensión exercise	10	23.1±4.1	
Cadore <i>et al.</i> (2014)	Concentric training	11	22.9 ± 7.8	Physically active
	Eccentric training	11	21.3 ± 3.3	
	Control	17	20.5 ± 2.3	
Coratella <i>et al.</i> (2015)	Isokinetic unilateral eccentric training	16	20 ± 1	Not mentioned
	Dynamic constant external resistance unilateral eccentric training	16		
	Control	10		
Guilhem <i>et al.</i> (2013)	Work-matched isoload eccentric resistance training	11	Not mentioned	Soccer players
	Isokinetic eccentric resistance training	10		
	Sprint training	10		
Mendiguchia <i>et al.</i> (2020)	Nordic hamstring exercise	12	26± 2.7	Moderately physically active
	Control	10		
Ribeiro-Alvares <i>et al.</i> (2018)	Nordic hamstring exercise	10	23.7 ± 3.3	Not mentioned
	Control	10	22,8 ± 4,2	
Sanz- López <i>et al.</i> (2017)	Eccentric Overload Training	10	22,8 ± 4,2	Not mentioned
	Control	10		
Sanz- López <i>et al.</i> (2016)	Eccentric Overload Training	10	19.9±1.2	Recreational physical Activity
	Control	10		
Seymore <i>et al.</i> (2017)	Nordic hamstring exercise	10	18.3±0.5	Recreational physical Activity
	Control	10	22,3 ± 4,2	
Timmins <i>et al.</i> (2016)	Concentric strength training	14	23.0 ± 2.8	Recreational physical Activity
	Eccentric strength training	14		
Marusic <i>et al.</i> (2020)	Control	16	24.2 ± 2.1	Recreational physical Activity
	Eccentric hamstring exercises in a lengthened position	18		
Marzilger <i>et al.</i> (2020)	Control	13	27.2 ± 4.1	Young active
	Four isokinetic eccentric training	28	27.1 ± 4.4	
Marzilger <i>et al.</i> (2019)	Control	14	27.2 ± 4.1	Young active
	Four isokinetic eccentric training	33	27.1 ± 4.4	
Timmins <i>et al.</i> (2021)	Hip-dominant flywheel exercise	13	22 ± 3	Soccer players
	Nordic hamstring exercise	14		
	Control	13		
Abián <i>et al.</i> (2020)	Single-leg decline squat exercise (3s)	12	20,8 ± 1,9	Moderately physically active
	Single-leg decline squat exercise (6s)	11	21.1 ± 1,2	
	Control	10	26.4 ± 4.1	
Presland <i>et al.</i> (2020)	Flywheel training with an additional eccentric-bias	10	29.2 ± 6.2	Recreational physical Activity
	Control	10	25 ± 3	
Franchi <i>et al.</i> (2014)	Concentric strength training	6	25 ± 3	Not mentioned
	Eccentric strength training	6		

Fourteen studies evaluated the effects of eccentric training on FL. Significant increases in FL were found in 12 studies, six of which (Bourne *et al.*; Mendiguchia *et al.*; Ribeiro-Alvares *et al.*; Timmins *et al.*, 2016, 2021; Marusic *et al.*) performed NHE strength training, five performed isokinetic eccentric strength training (Franchi *et al.*; Cadore *et al.*; Coratella *et al.*; Marzilger *et al.*, 2019, 2020) and one completed flywheel training with an additional eccentric bias (Presland *et al.*).

Eleven studies evaluated the effects of eccentric training on MT. Only in one study, no change in MT was found after flywheel training with an additional eccentric bias (Presland *et al.*). In five studies, there was a significant

increase in MT compared to baseline and after performing eccentric training with heel drop exercise (Alonso-Fernandez *et al.*), unilateral isokinetic eccentric training (Cadore *et al.*; Coratella *et al.*), work-matched isoload eccentric resistance training (Guilhem *et al.*), and unilateral constant external resistance dynamic eccentric training (Coratella *et al.*). Only one study assessed muscle quality through echo intensity (Cadore *et al.*), where they found a significant increase in muscle quality, decreasing its echo intensity. PSCA was evaluated in two studies (Sanz-Lopez *et al.*, 2017; Marzilger *et al.*, 2020), showing an increase in this parameter. The exercise modalities, the muscles tested and their effects on muscle architecture are summarized in Table III.

Table II. Methodological quality of the studies included (Y= yes; N= no).

Studies	Items											Score (/10)
	1	2	3	4	5	6	7	8	9	10	11	
Alonso- Fernández <i>et al.</i> (2020)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Bourne <i>et al.</i> (2017)	N	Y	N	N	N	N	Y	N	N	Y	Y	4/10
Cadore <i>et al.</i> (2014)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Coratella <i>et al.</i> (2015)	Y	Y	Y	Y	N	N	Y	Y	Y	Y	Y	8/10
Guilhem <i>et al.</i> (2013)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Mendiguchia <i>et al.</i> (2020)	Y	Y	N	N	N	N	Y	Y	Y	Y	Y	6/10
Ribeiro-Alvares <i>et al.</i> (2018)	Y	Y	N	Y	N	N	N	Y	Y	Y	Y	6/10
Sanz- López <i>et al.</i> (2017)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Sanz- López <i>et al.</i> 2016	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Seymore <i>et al.</i> (2017)	Y	Y	N	Y	N	N	N	Y	Y	Y	Y	6/10
Timmins <i>et al.</i> (2016)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Marusic <i>et al.</i> (2020)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Marzilger <i>et al.</i> (2020)	N	Y	N	Y	N	N	N	Y	Y	Y	Y	6/10
Marzilger <i>et al.</i> (2019)	Y	Y	N	Y	N	N	N	Y	Y	Y	Y	6/10
Timmins <i>et al.</i> (2021)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Abián <i>et al.</i> (2020)	Y	Y	N	Y	N	N	N	Y	Y	Y	Y	6/10
Presland <i>et al.</i> (2020)	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7/10
Franchi <i>et al.</i> (2014)	Y	N	N	Y	N	N	N	Y	Y	Y	Y	5/10

Items considered for rating: 1. Eligibility criteria were specified (This item is not used to calculate the PEDro score.); 2. Subjects were randomly allocated to groups (in a crossover study, subjects were randomly allocated an order in which treatments were received); 3. Allocation was concealed; 4. The groups were similar at baseline regarding the most important prognostic indicators; 5. There was blinding of all subjects; 6. There was blinding of all therapists who administered the therapy; 7. There was blinding of all assessors who measured at least one key outcome; 8. Measures of at least one key outcome were obtained from more than 85% of the subjects initially allocated to groups; 9. All subjects for whom outcome measures were available received the treatment or control condition as allocated or, where this was not the case, data for at least one key outcome was analysed by “intention to treat”; 10. The results of between-group statistical comparisons are reported for at least one key outcome; 11. The study provides both point measures and measures of variability for at least one key outcome.

## DISCUSSION

The present study aimed to evaluate the effects of eccentric training on muscle architecture. The parameters most frequently evaluated in the studies consulted were PA, FL, and MT. These were assessed mainly in lower limb muscles such as BF<sub>lh</sub>, VL, MG, and LG. Eccentric training for at least four weeks generates adaptations in these parameters, mainly by increasing MT with FL and decreasing PA, determining muscle function. These results provide evidence on the effects of eccentric training on muscle architecture, which could be helpful to prevent injuries and favor muscle recovery processes.

NHE was the most used eccentric training in this review. This type of training targeted the hamstring muscles, being widely used in the literature (Llurda-Almuzara *et al.*, 2021), perhaps because it is easy to perform and reproduce, as it requires only a partner and no dynamometric equipment is necessary (Gerard *et al.*), is efficient in reducing injury risk (Al Attar *et al.*, 2017), increasing eccentric strength (Ishøi *et al.*, 2018) and potentially modifying muscle architecture by increasing fascicle length (Medeiros *et al.*,

2020). The latter is repeated in all the studies that used NHE for this review, showing a significant increase in FL after performing their protocols, except as described by Seymore *et al.* who found no significant changes, possibly due to the joint positions assumed during training. A different situation is seen when reviewing the effects of this exercise on PA, showing a decrease in all the studies, except for those described by Seymore *et al.*, where there was an increase in PA without significant changes. Concerning MT, this exercise caused a variable increase in all the studies used.

Concerning the other eccentric training modalities included in this review, the second modality used was eccentric isokinetic exercises using a technological tool to control angular velocity. It has already been possible to determine the effects of isokinetic eccentric exercise on quadriceps muscle mass and strength in a population before or after ACL reconstruction or meniscectomy (Zhang *et al.*, 2017; Vidmar *et al.*, 2019, 2020), its effect on neural adaptations has also been seen (Barrué-Belou *et al.*, 2016). The results of this review support its use as a potential way

Table III. Exercise interventions.

Study	Type of exercise	weeks	Procedure	Assessed parameters	Muscle	FL	MT	PA	OM	PSCA or CSA
Alonso-Ferrández <i>et al.</i> (2020)	Eccentric training with heel drop exercise	8	2 sets x 10-12 repetitions (week 1) 3x/week, 3 x 12-15 (week 2) 3x/week, 3 sets x 15 repetitions (week 3 and 4) 4x/week, 4 sets x 15 repetitions (+2kg load) (week 5) 4x/week, 4 sets x 15 repetitions (week 6) 4x/week, 4 sets x 15 repetitions (week 7) 5x/week, 4 sets x 10-12 repetitions (week 8) 5x/week, 4 sets x 10-12 repetitions (week 8) 5x/week, 2-6 sets x 5-10 repetitions, rest 3 min between sets, 2x/week (alternate legs)	FL; MT; PA	MG; LG	**	**	**	N/A	N/A
Bourne <i>et al.</i> (2017)	NHE	10	2-6 sets x 5-10 repetitions, rest 3 min between sets, 2x/week (alternate legs)	FL	BFlh	##	N/A	N/A	N/A	N/A
Cadore <i>et al.</i> (2014)	Iso kinetic unilateral Eccentric training	6	2 sets x 8 repetitions, 2x/ weeks	MT; QM	VL	N/A	**	N/A	**	N/A
Coratella <i>et al.</i> (2015)	Iso kinetic unilateral eccentric training	8	5 sets x 8 repetitions, 2 x/week	FL; MT; PA	VL	**	**	-	N/A	N/A
	Dynamic constant external resistance unilateral eccentric training	8	5 sets x 8 repetitions, 2 x/week	FL; MT; PA	VL	**	**	-	N/A	N/A
Guilhem <i>et al.</i> (2013)	Work-matched iso load eccentric resistance training	9	3 sets x 8 repetitions (week 1) 4 sets x 8 repetitions (week 2-3) 5 sets x 8 repetitions (week 4, 6-9)	FL; MT; PA	VL	**	**	**	N/A	N/A
	Iso kinetic eccentric resistance training	9	3 sets x 8 repetitions (week 1) 4 sets x 8 repetitions (week 2-3) 5 sets x 8 repetitions (week 4, 6-9)	FL; MT; PA	VL	**	**	**	N/A	N/A
Mendiguchia <i>et al.</i> (2020)	NHE	6	3 sets of 12, 10 and 8 repetitions respectively, 1x/week	FL; MT; PA	BFlh	**	**	**	N/A	N/A
Ribeiro-Alvares <i>et al.</i> (2018)	NHE	4	3 sets 3-6-10 repetitions, 1-min rest between sets, 2x/week	FL; MT; PA	BFlh	**	**	**	N/A	N/A
Sanz-López <i>et al.</i> (2017)	Eccentric Overhead Training	6	4 sets x 7 repetitions with 2 minutes of rest	PA	MG	N/A	N/A	**	N/A	** (PCSA)
Sanz-López <i>et al.</i> (2016)	Eccentric Overhead Training	6	5 sets x 7 repetitions with 2 minutes of rest	PA	VL	N/A	N/A	**	N/A	N/A
Seymore <i>et al.</i> (2017)	NHE	6	2 sets x 5 repetitions (week 1) 1x/week, 2 sets x 6 repetitions (week 2) 2x/week, 3 sets x 6-8 repetitions (week 3) 3x/week, 3 sets x 8-10 repetitions (week 4) 3x/week, 3 sets x 12-10-8 repetitions (week 5-6)	FL; PA; PSCA	BFlh	N/A	N/A	-	N/A	**
Timmins <i>et al.</i> (2016)	NHE	6	4 sets x 6 reps (week 1) 2x/week, 4 sets x 6 reps (week 2) 3x/week, 5 sets x 6 reps (week 3) 3x/week, 5 sets x 8 reps (week 4) 3x/week, 6 sets x 6 reps (week 5) 3x/week, 6 sets x 8 reps (week 6) 3x/week.	FL; MT; PA	BFlh	**	**	**	N/A	N/A
Manusi <i>et al.</i> (2020)	NHE in a lengthened position	6	2 sets x 5 repetitions (week 1), 2 sets x 6 repetitions (week 2), 3 sets x 6 repetitions (week 3), 3 sets x 8 repetitions (week 4-6) / all 2x/week	FL; MT; PA	BFLH	#	-	#	N/A	N/A
Marzilger <i>et al.</i> (2020)	Four isokinetic eccentric training (45°/s, 120°/s, 210°/s and 300°/s)	11	5 sets x 3, 8, 14 and 20 repetitions per-training set respectively (45°/s, 120°/s, 210°/s and 300°/s), 3x/week	FL; PA; PSCA	VL	**	N/A	-	N/A	(PCSA) p210/p300
Marzilger <i>et al.</i> (2019)	Four isokinetic eccentric training (45°/s, 120°/s, 210°/s and 300°/s)	11	5 sets x 3, 8, 14 and 20 repetitions per-training set respectively (45°/s, 120°/s, 210°/s and 300°/s), 3x/week	PSCA	VL	N/A	N/A	N/A	N/A	(CSA)
Timmins <i>et al.</i> (2021)	NHE	39	2 sessions per week after field training	FL; MT; PA	BFlh	**	-	**	N/A	N/A
Abián <i>et al.</i> (2020)	Single-leg decline squat exercise (3s)	6	3 sets x 8 repetitions, 3x/ week	FL; MT; PA	VL	-	#	##	N/A	N/A
	Single-leg decline squat exercise (6s)	6	3 sets x 8 repetitions, 3x/ week	FL; MT; PA	BFlh	-	**	-	N/A	N/A
Presland <i>et al.</i> (2020)	Flyw heel training with an additional eccentric-bias	6	4 sets x 6 repetitions (week 1 and 2), 5 sets x 6 reps (week 3), 5 sets x 8 reps (week 4), 6 sets x 8 repetitions (week 5), 5 sets x 6 reps (week 6)	FL; MT; PA	BFlh	##	NC	-	N/A	N/A
Franchi <i>et al.</i> (2014)	Iso kinetic eccentric training	10	4 sets x 8-10 repetitions, 3x/ (week)	FL; PA	VL	##	N/A	#	N/A	N/A

FL= fascicle length; MT= muscle thickness; PA=pennation angle; QM= muscle quality BFlh= biceps femoris long head; MG= medial gastrocnemius; LG= lateral gastrocnemius; VL= vastus lateralis; CSA= cross section; PCSA= physiological cross section area; NC: no changes. # = Significant changes vs. control group \*\* = Significant changes vs. baseline

of modifying muscle architecture parameters generating an increase in FL and MT, and a decrease in PA (Guilhem *et al.*; Marzilger *et al.*, 2020; Abián *et al.*; Coratella *et al.*; Cadore *et al.*).

On the other hand, when reviewing the effects of eccentric exercise on the architecture of different muscles, only studies that measured muscle architecture parameters in the lower limb muscles were found in this review. These were BFlh, VL, GM, and GL, respectively. Usually, the assessment of muscle architecture parameters is measured using a B-mode ultrasound machine, which has already been validated in lower limb muscles for this type of measurement (Van Hooren *et al.*, 2020; Turton *et al.*, 2019), which could explain the preference in choosing these muscles to assess. However, it is advisable to investigate the effects of eccentric exercise on upper limb muscles.

It is known that muscles with higher FL have a greater advantage in speed movements (Lieber & Ward), on the other hand, muscles with higher PA are muscles that can generate more force (Blazevich & Sharp). In this context, and according to the results obtained in this review, eccentric training could favor muscle function in speed situations, however, it

could disfavor it in strength situations. For this, it is important to distinguish the type of strength we refer to, so it is suggested to include the association with strength in future studies that attempt to resolve this research question. This review only considered the effects of eccentric training on architectural parameters, but not its association with strength or speed.

The population's average age included in the studies of this review ranges between 18 and 29 years of age. This prevents us from having a clearer picture of the general population; there is a lack of studies that evaluate the effects of eccentric training in the elderly population since the changes produced in skeletal muscle function with age are indisputable (Tieland *et al.*, 2018)

## CONCLUSION

The results of this review show that there are adaptations in muscle architecture to eccentric training of at least four weeks duration. This will allow health and sports professionals to understand in greater depth the adaptations of skeletal muscle to this type of exercise, and in this way, prevent injuries, favor the muscle repair process, and improve sports training. However, although this review includes a universe of more than 500 participants, the adaptations in muscle architecture in the upper limb musculature or the population of older adults and their association with muscle strength are still not known with certainty. In this context, it is suggested to conduct studies that include this population and the upper limb muscles.

## ACKNOWLEDGMENTS

This paper will be part of Rodrigo Lizama-Perez Doctoral Thesis performed in the Biomedicine Doctorate Program of the University of Granada, Spain.

---

**LIZAMA-PÉREZ, R.; CHIROSA-RIOS, I.; CHIROSA-RIOS, L.; OLAVE, E.; FERRAGUT, C.; VILA, H. & JEREZ-MAYORGA, D.** Efectos del ejercicio excéntrico en la arquitectura muscular en adultos: Una revisión sistemática. *Int. J. Morphol.*, 40(2):425-432, 2022.

**RESUMEN:** El propósito de esta revisión sistemática fue determinar los efectos del entrenamiento excéntrico sobre la arquitectura muscular en la población adulta. Se siguieron las recomendaciones del Items de referencia para publicar Revisiones Sistemáticas y Metaanálisis (PRISMA) utilizando palabras clave aso-

ciadas con la arquitectura muscular y el entrenamiento excéntrico en cuatro bases de datos: PubMed, Scopus, SPORTDiscus y Web of Science. La calidad metodológica se evaluó mediante la escala PEDro. Se encontró un total de 1260 artículos, del los cuales, 18 fueron incluidos en esta revisión. Los parámetros más frecuentemente evaluados en los estudios fueron el ángulo de penación (AP), la longitud del fascículo (LF) y el grosor muscular (Gm). Estos fueron evaluados principalmente en músculos de los miembros inferiores como la cabeza larga del bíceps femoral (CLBf), el vasto lateral (VL), el gastrocnemio medial (GM) y el gastrocnemio lateral (GL), respectivamente. El entrenamiento excéntrico durante al menos cuatro semanas genera adaptaciones en estos parámetros, principalmente aumentando el GM con la LF y disminuyendo el AP, determinando de esta manera la función muscular. Estos resultados aportan evidencias sobre los efectos del entrenamiento excéntrico en la arquitectura muscular, que podrían ser útiles para prevenir lesiones y favorecer los procesos de recuperación muscular.

**PALABRAS CLAVE:** Arquitectura muscular; Entrenamiento excéntrico; Potencia muscular; Longitud del fascículo; Angulo de penación; Grosor muscular; Calidad muscular.

## REFERENCES

- Abián, P.; Martínez F.; Jiménez F. & Abián-Vicén J. Effects of eccentric single-leg decline squat exercise on the morphological and structural properties of the vastus lateralis and patellar tendon. *Int. J. Environ. Res. Public Health.*, 17(24):1-12, 2020.
- Al Attar, W.S.A.; Soomro, N.; Sinclair, P.J.; Pappas, E. & Sanders, R.H. Effect of injury prevention programs that include the nordic hamstring exercise on hamstring injury rates in soccer players: a systematic review and meta-analysis. *Sport Med.*, 47(5):907-16, 2017.
- Alonso-Fernandez, D.; Taboada-Iglesias, Y.; García-Remeseiro, T. & Gutiérrez-Sánchez, Á. Effects of the functional heel drop exercise on the muscle architecture of the gastrocnemius. *J. Sport. Rehabil.*, 29(8):1053-9, 2020.
- Barrué-Belou, S.; Amarantini, D.; Marque, P. & Duclay, J. Neural adaptations to submaximal isokinetic eccentric strength training. *Eur. J. Appl. Physiol.*, 116(5):1021-30, 2016.
- Blazevich, A. J. & Sharp, N. C. C. Understanding muscle architectural adaptation: macro- and micro-level research. *Cells Tissues Organs*, 181(1):1-10, 2005.
- Bourne, M. N.; Duhig, S. J.; Timmins, R. G.; Williams, M. D, Opar, D. A.; Al Najjar, A.; Kerr, G. K. & Shield, A. J. Impact of the Nordic hamstring and hip extension exercises on hamstring architecture and morphology: Implications for injury prevention. *Br. J. Sports. Med.*, 51(5):469-77, 2017.
- Cadore, E. L.; González-Izal, M.; Pallarés, J.G.; Rodríguez-Falces, J.; Häkkinen, K.; Kraemer, W. J.; Pinto, R. S. & Izquierdo, M. Muscle conduction velocity, strength, neural activity, and morphological changes after eccentric and concentric training. *Scand. J. Med. Sci. Sports.*, 24(5):e343-52, 2014.
- Cashin, A. G. & McAuley, J. H. Clinimetrics: Physiotherapy Evidence Database (PEDro) Scale. *J. Physiother.*, 66(1):59, 2020.
- Coratella, G.; Milanese, C. & Schena, F. Unilateral eccentric resistance training: A direct comparison between isokinetic and dynamic constant external resistance modalities. *Eur. J. Sport. Sci.*, 15 (8):720-6, 2015.
- Fragala, M. S.; Fukuda, D. H.; Stout, J. R.; Townsend, J. R.; Emerson, N. S.; Boone, C. H.; Beyer, K. S.; Oliveira, L. P. & Hoffman, J. R. Muscle

- quality index improves with resistance exercise training in older adults. *Exp. Gerontol.*, 53:1-6, 2014.
- Fragala, M. S.; Kenny, A. M. & Kuchel, G. A. Muscle quality in aging: a multi-dimensional approach to muscle functioning with applications for treatment. *Sport Med.*, 45(5):641-58, 2015.
- Franchi, M. V.; Atherton, P. J.; Reeves, N. D.; Flück, M.; Williams, J.; Mitchell, W. K.; Selby, A.; Beltran Valls, R. M. & Narici, M. V. Architectural, functional and molecular responses to concentric and eccentric loading in human skeletal muscle. *Acta Physiol. (Oxf)*, 210(3):642-54, 2014.
- Gerard, R.; Gojon, L.; Declève, P. & Van Cant, J. The effects of eccentric training on biceps femoris architecture and strength: a systematic review with meta-analysis. *J. Athl. Train.*, 55(5):501-14, 2020.
- Guilhem, G.; Cornu, C.; Maffiuletti, N. A. & Guével, A. Neuromuscular adaptations to isoload versus isokinetic eccentric resistance training. *Med. Sci. Sports Exerc.*, 45(2):326-35, 2013.
- Ishoi, L.; Hölmich, P.; Aagaard, P.; Thorborg, K.; Bandholm, T. & Serner, A. Effects of the Nordic Hamstring exercise on sprint capacity in male football players: a randomized controlled trial. *J. Sports. Sci.*, 36(14):1663-72, 2018.
- Jerez-Mayorga, D.; Delgado-Floody, P.; Intelangelo, L.; Campos-Jara, C.; Arias-Poblete, L.; García-Verazaluce, J.; García-Ramos, A.; & Chiroso, L. J. Behavior of the muscle quality index and isometric strength in elderly women. *Physiol. Behav.*, 227:113145, 2020.
- Lieber, R. L. & Fridén, J. Functional and clinical significance. *Muscle Nerve*, 23:1647-66, 2000.
- Lieber, R. L. & Ward, S. R. Skeletal muscle design to meet functional demands. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 366(1570):1466-76, 2011.
- Llurda-Almuzara, L.; Labata-Lezaun, N.; López-de-Celis, C.; Aiguadé-Aiguadé, R.; Romaní-Sánchez, S.; Rodríguez-Sanz, J.; Fernández-de-Las-Peñas, C. & Pérez-Bellmunt, A. Biceps femoris activation during hamstring strength exercises: a systematic review. *Int. J. Environ. Res. Public Health*, 18(16):8733, 2021.
- Marušić, J.; Vatovec, R.; Marković, G. & Šarabon, N. Effects of eccentric training at long-muscle length on architectural and functional characteristics of the hamstrings. *Scand. J. Med. Sci. Sport.*, 30(11):2130-42, 2020.
- Marzilger, R.; Bohm, S.; Mersmann, F. & Arampatzis, A. Modulation of physiological cross-sectional area and fascicle length of vastus lateralis muscle in response to eccentric exercise. *J. Biomech.*, 111:110016, 2020.
- Marzilger, R.; Bohm, S.; Mersmann, F. & Arampatzis, A. Effects of lengthening velocity during eccentric training on vastus lateralis muscle hypertrophy. *Front. Physiol.*, 10:957, 2019.
- Medeiros, D. M.; Marchiori, C. & Baroni, B. M. Effect of nordic hamstring exercise training on knee flexors eccentric strength and fascicle length: a systematic review and meta-analysis. *J. Sport. Rehabil.*, 12;30(3):482-91, 2020.
- Mendiguchia, J.; Conceição, F.; Edouard, P.; Fonseca, M.; Pereira, R.; Lopes, H.; Morin, J. B. & Jiménez-Reyes, P. Sprint versus isolated eccentric training: comparative effects on hamstring architecture and performance in soccer players. *PLoS One*, 15(2):e0228283, 2020.
- Naimo, M. A.; Varanoske, A. N.; Hughes, J. M. & Pasiakos, S. M. Skeletal muscle quality: a biomarker for assessing physical performance capabilities in young populations. *Front. Physiol.*, 12:706699, 2021.
- Narici, M.; Franchi, M. & Maganaris, C. Muscle structural assembly and functional consequences. *J. Exp. Biol.*, 219(Pt. 2):276-84, 2016.
- Ouzzani, M.; Hammady, H.; Fedorowicz, Z. & Elmagarmid, A. Rayyan—a web and mobile app for systematic reviews. *Syst. Rev.*, 5:210, 2016.
- Presland, J. D.; Opar, D. A.; Williams, M. D.; Hickey, J. T.; Maniar, N.; Lee Dow, C.; Bourne, M. & Timmins, R. Hamstring strength and architectural adaptations following inertial flywheel resistance training. *J. Sci. Med. Sport.*, 23(11):1093-9, 2020.
- Ribeiro-Alvares, J. B.; Marques, V. B.; Vaz, M. A. & Baroni, B. M. Four weeks of nordic hamstring exercise reduce muscle injury risk factors in young adults. *J. Strength. Cond. Res.*, 32(5):1254-62, 2018.
- Sanz-López, F.; Berzosa Sánchez, C.; Hita-Contreras, F.; Cruz-Díaz, D. & Martínez-Amat, A. Ultrasound changes in Achilles tendon and gastrocnemius medialis muscle on squat eccentric overload and running performance. *J. Strength Cond. Res.*, 30(7):2010-8, 2016.
- Sanz-López, F.; Berzosa, C.; Hita-Contreras, F. & Martínez-Amat, A. Effects of eccentric overload training on patellar tendon and vastus lateralis in three days of consecutive running. *Knee*, 24(3):570-9, 2017.
- Seymore, K. D.; Doherty, Z. J.; DeVita, P.; Rider, P. M. & Kulas, A. S. The effect of Nordic hamstring strength training on muscle architecture, stiffness, and strength. *Eur. J. Appl. Physiol.*, 117(5):943-53, 2017.
- Suchomel, T. J.; Nimphius, S. & Stone, M. H. The importance of muscular strength in athletic performance. *Sports Med.*, 46(10):1419-49, 2016.
- Tieland, M.; Trouwborst, I. & Clark, B. C. Skeletal muscle performance and ageing. *J. Cachexia Sarcopenia Muscle*, 9(1):3-19, 2018.
- Timmins, R. G.; Filippoulos, D.; Nguyen, V.; Giannakis, J.; Ruddy, J. D.; Hickey, J. T.; Maniar, N. & Opar, D. Sprinting, strength, and architectural adaptations following hamstring training in Australian footballers. *Scand. J. Med. Sci. Sport*, 31(6):1276-89, 2021.
- Timmins, R. G.; Ruddy, J. D.; Presland, J.; Maniar, N.; Shield, A. J.; Williams, M. D. & Opar, D. Architectural changes of the biceps femoris long head after concentric or eccentric training. *Med. Sci. Sports Exerc.*, 48(3):499-508, 2016.
- Turton, P.; Hay, R. & Welters, I. Assessment of peripheral muscle thickness and architecture in healthy volunteers using hand-held ultrasound devices; a comparison study with standard ultrasound. *BMC Med. Imaging*, 19(1):69, 2019.
- Van Hooren, B.; Teratsias, P. & Hodson-Tole, E. F. Ultrasound imaging to assess skeletal muscle architecture during movements: a systematic review of methods, reliability, and challenges. *J. Appl. Physiol.* (1985), 128(4):978-99, 2020.
- Vidmar, M. F.; Baroni, B. M.; Michelin, A. F.; Mezzomo, M.; Lugokenski, R.; Pimentel, G. L. & Silva, M. F. Isokinetic eccentric training is more effective than constant load eccentric training for quadriceps rehabilitation following anterior cruciate ligament reconstruction: a randomized controlled trial. *Braz. J. Phys. Ther.*, 24(5):424-32, 2020.
- Vidmar, M. F.; Baroni, B. M.; Michelin, A. F.; Mezzomo, M.; Lugokenski, R.; Pimentel, G. L. & Silva, M. F. Isokinetic eccentric training is more effective than constant load eccentric training on the quadriceps rehabilitation following partial meniscectomy: A randomized clinical trial. *Phys. Ther. Sport*, 39:120-5, 2019.
- Zhang, X. Y.; Huang, H. S.; Yang, J.; Yu, Y. Y.; Zhang, D. X. & Chang, C. Q. Effects of preoperative isokinetic eccentric training and whey protein isolate supplement on quadriceps strength and knee function in patients with anterior cruciate ligament rupture. *Zhongguo. Yi Xue Ke Xue Yuan Xue Bao*, 39(6):792-9, 2017.

Corresponding author:  
Daniel Jerez-Mayorga  
PhD in Biomedicine  
Fernandez Concha 700  
Las Condes  
Santiago  
CHILE

E-mail: daniel.jerez@unab.cl

ORCID ID: 0000-0002-6878-8004